Page 1 of 2

MUSURCE INEVLINE/AIDSLINE





THE POTE SHIP **Numbers Here** 

Search

Medscape Search | MEDLINE Search | Druginfo Search

My Homepage | Log Out BEST AVAILABLE COPY

October 1, 2004

Order Full Text

Add to Clipboard

Olfactory threshold shift following controlled 7-hour exposure to toluene and/or xylene.

Neurotoxicology 1992 Spring;13(1);211-5 (ISSN: 0161-813X)

Mergler D: Beauvais B

Centre pour l'etude des interactions biologiques entre la sante et l'environnement (CINBIOSE), Universite du Quebec a Montreal, Canada.

The present study was undertaken to examine olfactory perception threshold (OPT) shift following exposure of healthy subjects to toluene and/or xylene in an inhalation chamber. Five volunteers were exposed to 50 ppm toluene, 40 ppm xylene or an additive mixture of the two, for a period of 7 consecutive hr in an inhalation chamber. A Latin square design was used and subjects were exposed over 3 consecutive days/week, with an 11-day interval between each 3-day session. Olfactory perception thresholds, measured in decismets (ds), were ascertained for both toluene and PM-carbinol\_contained in 100 ml bottles with senally increasing concentrations (Offacto-Lab Kits # 191 & 11). Test administration was based on the forced choice method. Analysis of variance of pre-exposure OPTs indicated that for both toluene and PM-carbinol, significant differences were observed between individuals (pless than 0.05), but not between days or weeks. Measurements, made immediately following exposures revealed a significant six-fold increase in OPT for toluene (median: 15 ds), while PM-carbinol OPT remained stable. Individual differences were observed, but there was no effect of type of exposure, day, week, or interactions. OPT for toluene, determined at intervals following cessation of exposure, indicated a return to pre-exposure values at a mean rate of 6.8 ds/hr. The findings of this study suggest that there is a substantial offactory threshold shift during a 7-hr period, specific to a particular solvent or family of solvents. Receptorspecific saturation is proposed as the underlying mechanism

Major Subject Heading(s)	Minor Subject Heading(s)	CAS Registry / EC Numbers
<ul> <li>Sensory Thresholds [drug effects]</li> <li>Smell [drug effects]</li> <li>Toluene [administration &amp; dosage]</li> <li>Xylenes [administration &amp; dosage]</li> </ul>	<ul> <li>Administration,</li> <li>Inhalation</li> <li>Adult</li> <li>Middle Aged</li> <li>Reference Values</li> <li>Time Factors</li> </ul>	0 (Xylenes)     108-88-3 (Toluene)

"I didn't think 5 Pounds matterest We Have the Solution STARTHER Lilida't tainic 5 Pounds mattered We Havs the Solution START HERE Visit the

WebMB Health

Coasumer Site.

CLICK HERE!

Indexing Check Tags: Human

Language: English

MEDLINE Indexing Date: 199209 Publication Type: Owner: NLM Publication Type: Journal Article PreMedline Identifier: 0001508422

Journal Code: IM

Add to Clipboard



SEARCH

Medscape

Druginfo

**MEDLINE** 

Advanced Search



About Medscape - Privacy & Ethics - Terms of Use - Help - WebMD Health